

RESIDUES FOR AKIZUKI'S ONE-DIMENSIONAL LOCAL DOMAIN

I-CHIAU HUANG AND JAN-LI LIN

ABSTRACT. For a one-dimensional local domain C_M constructed by Akizuki, we find residue maps which give rise to a local duality. The completion of C_M is described using these residue maps.

Injective hulls of a given module are all isomorphic. For this reason, people often speak of *the* injective hull to indicate its “*uniqueness*”. However, isomorphisms between these injective hulls are not canonical. In fact, they are a part of the structure of the given module. For instance, a local duality for a power series ring [2, (5.9)] is interpreted as an isomorphism between two injective hulls - one given by local cohomology and another by continuous homomorphisms. This isomorphism is induced by a residue map, which was not observed from the viewpoint of “*uniqueness*” of injective hulls. In this article, our philosophy is taken up again by a Noetherian local ring C_M constructed by Akizuki [1]. Although C_M behaves beyond geometric expectation, we can still define certain maps, which give rise to a local duality as an identification of local cohomology classes and continuous homomorphisms. These maps, also called residue maps, determine all endomorphisms of an injective hull of the residue field of C_M . So we are able to describe the completion of C_M .

We recall Akizuki's construction. Let A be a discrete valuation ring with the maximal ideal $\mathfrak{m} = tA$, let $\hat{A}$ be its completion, and let K (resp. $\hat{K}$) be the quotient field of A (resp. $\hat{A}$). Assume that there is an element

$$z = a_0 + a_1 t^{n_1} + a_2 t^{n_2} + \cdots \in \hat{A} \quad (a_i \in A \setminus \mathfrak{m})$$

transcendental over A with the condition

$$n_r \geq 2n_{r-1} + 2 \quad (r \geq 1)$$

on exponents, where $n_0 = 0$. Let

$$z_r = a_r + a_{r+1} t^{n_{r+1} - n_r} + \cdots \quad (r \geq 0)$$

and

$$C = A[t(z_0 - a_0), \{(z_i - a_i)^2\}_{i=0}^\infty].$$

C_M is defined to be the localization of C at the maximal ideal M generated by t and $t(z_0 - a_0)$. Akizuki [1] showed that C_M is a one-dimensional Noetherian local domain, whose normalization is not a finite C_M -module.

The quotient field of C_M is $K(z)$, which equals $(C_M)_t$ as t is a system of parameter of C_M . We can use the exact sequence

$$(1) \quad 0 \rightarrow C_M \xrightarrow{\text{localization}} (C_M)_t \rightarrow H_{MC_M}^1(C_M) \rightarrow 0$$

1991 *Mathematics Subject Classification.* Primary 13H10; Secondary 13D45.

Key words and phrases. Gorenstein ring, injective hull, local duality, residue.

to describe elements of the first local cohomology module $H_{MC_M}^1(C_M)$ of C_M supported at the maximal ideal of C_M : For $f \in C_M$ and $n > 0$, the generalized fraction

$$\begin{bmatrix} f \\ t^n \end{bmatrix}_{C_M}$$

is defined as the image of f/t^n in $H_{MC_M}^1(C_M)$ under the map in (1). The description of top local cohomology modules by generalized fractions is essentially used in the concrete realizations of Grothendieck duality [2, 4, 5]. It can be applied to combinatorial analysis [3, 6]. See also [9] for an alternate treatment to generalized fractions.

For $f \in \hat{A}$ and $n > 0$, we denote by

$$(2) \quad \begin{bmatrix} f \\ t^n \end{bmatrix}_A$$

the image of f/t^n in $\hat{K}/\hat{A}$ under the map in the exact sequence

$$0 \rightarrow \hat{A} \xrightarrow{\text{inclusion}} \hat{K} \rightarrow \hat{K}/\hat{A} \rightarrow 0.$$

We remark that there is a canonical isomorphism between $\hat{K}/\hat{A}$ and $H_{\mathfrak{m}\hat{A}}^1(\hat{A})$, with which the representation (2) of elements in $\hat{K}/\hat{A}$ and the representation of elements in $H_{\mathfrak{m}\hat{A}}^1(\hat{A})$ by generalized fractions agree.

Vanishing of elements in $\hat{K}/\hat{A}$ and $H_{MC_M}^1(C_M)$ can be described in terms of ideal membership:

$$\begin{bmatrix} f \\ t^n \end{bmatrix}_A = 0 \quad \left(\text{resp. } \begin{bmatrix} f \\ t^n \end{bmatrix}_{C_M} = 0 \right)$$

if and only if f is contained in the ideal of $\hat{A}$ (resp. C_M) generated by t^n . For instance,

$$\begin{bmatrix} t(z_0 - a_0) \\ t \end{bmatrix}_A = 0$$

but

$$\begin{bmatrix} t(z_0 - a_0) \\ t \end{bmatrix}_{C_M} \neq 0.$$

The canonical map $K/A \rightarrow \hat{K}/\hat{A}$ is an isomorphism. We use the notation in (2) to represent elements in K/A under this isomorphism.

Lemma 1. *Every element of $H_{MC_M}^1(C_M)$ can be written as*

$$\begin{bmatrix} X + Yt(z_0 - a_0) \\ t^n \end{bmatrix}_{C_M}$$

for some $X, Y \in A$, $n \geq 0$.

Proof. It suffices to show that, for any specified $n > 0$, any element $f \in C_M$ can be written as

$$(3) \quad f = X + Yt(z_0 - a_0) + t^n Z$$

with $X, Y \in A$ and $Z \in C_M$. Write f as

$$f = \frac{f_1}{1 - f_2}$$

for some $f_1 \in C$ and $f_2 \in M$. Then

$$f - f_1(1 + f_2 + f_2^2 + \cdots + f_2^n) = \frac{f_1 f_2^{n+1}}{1 - f_2} \in M^{n+1}C_M.$$

Since $M^{n+1}C_M \subset t^n C_M$,

$$f - f_1(1 + f_2 + f_2^2 + \cdots + f_2^n) = t^n Z_2$$

for some $Z_2 \in C_M$. By [8, p. 140, Section 9.5, equation (5)], there exist $X, Y \in A$ and $Z_1 \in C$ such that

$$f_1(1 + f_2 + f_2^2 + \cdots + f_2^n) = X + Yt(z_0 - a_0) + t^n Z_1.$$

The representation

$$f = X + Yt(z_0 - a_0) + t^n(Z_1 + Z_2)$$

is of the required form. $\square$

Lemma 2. *If $X, Y \in A$ and*

$$\begin{bmatrix} X + Yt(z_0 - a_0) \\ t^n \end{bmatrix}_{C_M} = 0,$$

then $X, Y \in t^n A$.

Proof.

$$X + Yt(z_0 - a_0) = t^n Z$$

for some $Z \in C_M$. Write $X = t^\ell X_1$ and $Y = t^m Y_1$ with invertible $X_1, Y_1 \in A$. If $\ell \leq m$, then $n \leq \ell$, otherwise $X_1 = -t^{m-\ell} Y_1 t(z_0 - a_0) + t^{n-\ell} Z \in t\hat{A}$. If $\ell > m$, then $m \geq n$, otherwise $t(z_0 - a_0) = -t^{\ell-m} X_1 Y_1^{-1} + t^{n-m} Z Y_1^{-1} \in tC_M$. In either case, $X, Y \in t^n A$. $\square$

With these lemmas, we are able to define the following map for any $\sigma, \rho \in \hat{A}$.

Definition 3.

$$\text{res}_{\sigma, \rho}: H_{MC_M}^1(C_M) \rightarrow K/A$$

is defined to be the A -linear map given by

$$\begin{bmatrix} X + Yt(z_0 - a_0) \\ t^n \end{bmatrix}_{C_M} \mapsto \begin{bmatrix} X\sigma + Y\rho \\ t^n \end{bmatrix}_A.$$

Adopting the terminology of [2], we call $\text{res}_{\sigma, \rho}$ a residue map. Let

$$\text{Hom}_A^c(C_M, K/A) = \{\varphi \in \text{Hom}_A(C_M, K/A) \mid \varphi(M^n C_M) = 0 \text{ for some } n\}$$

be the C_M -module of continuous homomorphism. As a special case of J. Lipman's result [2, Proposition 3.4], $\text{Hom}_A^c(C_M, K/A)$ is an injective hull of the residue field of C_M . Note that $t^{n+1}C_M \subset M^{n+1}C_M \subset t^n C_M$. Hence a A -linear map $\varphi: C_M \rightarrow K/A$ is continuous if and only if $\varphi(t^n C_M) = 0$ for some n . Using the representation (3) of elements of C_M , we see that a continuous homomorphism is determined by an integer n with which $t^n C_M$ is in the kernel and by its values at 1 and $t(z_0 - a_0)$.

Definition 4.

$$\Phi_{\sigma, \rho}: H_{MC_M}^1(C_M) \rightarrow \text{Hom}_A^c(C_M, K/A)$$

is defined to be the C_M -linear map given by

$$\Phi_{\sigma, \rho}(\omega)(f) = \text{res}_{\sigma, \rho}(f\omega),$$

where $\omega \in H_{MC_M}^1(C_M)$ and $f \in C_M$.

Local Duality. *If ρ is invertible, then $\Phi_{\sigma,\rho}$ is an isomorphism.*

Proof. The inverse map of $\Phi_{\sigma,\rho}$ can be written down explicitly. Let

$$s_r := a_1 t^{n_1} + a_2 t^{n_2} + \cdots + a_r t^{n_r} \in A.$$

Then we have

$$t(z_0 - a_0) = t^{n_r+1}(z_r - a_r) + ts_r$$

and

$$(4) \quad t^2(z_0 - a_0)^2 + t^2 s_r^2 - 2ts_r t(z_0 - a_0) = t^{2n_r+2}(z_r - a_r)^2 \in t^r C_M.$$

Given $\varphi \in \text{Hom}_A^c(C_M, K/A)$ with $\varphi(t^r C_M) = 0$ and

$$\begin{cases} \varphi(1) = \begin{bmatrix} \alpha \\ t^r \end{bmatrix}_A \\ \varphi(t(z_0 - a_0)) = \begin{bmatrix} \beta \\ t^r \end{bmatrix}_A \end{cases},$$

the system of equations

$$\begin{cases} \begin{bmatrix} X\sigma + Y\rho \\ t^r \end{bmatrix}_A = \begin{bmatrix} \alpha \\ t^r \end{bmatrix}_A \\ \begin{bmatrix} X\rho - Yt^2 s_r^2 \sigma + 2Yts_r \rho \\ t^r \end{bmatrix}_A = \begin{bmatrix} \beta \\ t^r \end{bmatrix}_A \end{cases}$$

can be solved by choosing $X, Y \in A$ such that

$$\begin{cases} X - \frac{\alpha ts_r(\sigma ts_r - 2\rho) + \beta \rho}{(\rho - ts_r \sigma)^2} \in t^r \hat{A} \\ Y - \frac{\alpha \rho - \beta \sigma}{(\rho - ts_r \sigma)^2} \in t^r \hat{A}. \end{cases}$$

We define

$$\Phi^{-1}(\varphi) := \begin{bmatrix} X + Yt(z_0 - a_0) \\ t^r \end{bmatrix}_{C_M},$$

which is independent of the choices of r, α, β, X or Y . Then

$$\Phi_{\sigma,\rho}(\Phi^{-1}(\varphi))(1) = \begin{bmatrix} \alpha \\ t^r \end{bmatrix}_A$$

and

$$\begin{aligned} & \Phi_{\sigma,\rho}(\Phi^{-1}(\varphi))(t(z_0 - a_0)) \\ &= \text{res}_{\sigma,\rho} \left[\begin{bmatrix} Xt(z_0 - a_0) - Yt^2 s_r^2 + 2Yts_r t(z_0 - a_0) \\ t^r \end{bmatrix}_{C_M} \right] \quad (\text{by (4)}) \\ &= \begin{bmatrix} \beta \\ t^r \end{bmatrix}_A. \end{aligned}$$

Hence $\Phi_{\sigma,\rho}(\Phi^{-1}(\varphi)) = \varphi$. For any $\omega \in H_{MC_M}^1(C_M)$, it is also straightforward to check that $\Phi^{-1}(\Phi_{\sigma,\rho}(\omega)) = \omega$. So Φ^{-1} is indeed the inverse of $\Phi_{\sigma,\rho}$. $\square$

Corollary 5. C_M is Gorenstein.

Proof. $H_{MC_M}^1(C_M)$ is injective. Hence (1) is a finite injective resolution of C_M . $\square$

We remark that a Noetherian local ring R whose maximal ideal is generated by $1 + \text{depth } R$ elements is always Gorenstein [7, p. 163, Exercise 1].

Proposition 6. *Any C_M -linear map*

$$\Phi: H_{MC_M}^1(C_M) \rightarrow \text{Hom}_A^c(C_M, K/A)$$

equals $\Phi_{\sigma,\rho}$ for some $\sigma, \rho \in \hat{A}$.

Proof. For each n , there exist $\sigma_n, \rho_n \in A$ such that

$$\begin{cases} \begin{bmatrix} \sigma_n \\ t^n \end{bmatrix}_A = \Phi \left(\begin{bmatrix} 1 \\ t^n \end{bmatrix}_{C_M} \right) (1) \\ \begin{bmatrix} \rho_n \\ t^n \end{bmatrix}_A = \Phi \left(\begin{bmatrix} 1 \\ t^n \end{bmatrix}_{C_M} \right) (t(z_0 - a_0)). \end{cases}$$

Since $\sigma_n - \sigma_{n+1}$ and $\rho_n - \rho_{n+1}$ are contained in $t^n A$, the limits

$$\begin{cases} \sigma = \lim_{n \rightarrow \infty} \sigma_n \\ \rho = \lim_{n \rightarrow \infty} \rho_n \end{cases}$$

exist in $\hat{A}$. For any $X, Y \in A$,

$$\Phi \left(\begin{bmatrix} X + Yt(z_0 - a_0) \\ t^n \end{bmatrix}_{C_M} \right) (1) = \begin{bmatrix} X\sigma_n + Y\rho_n \\ t^n \end{bmatrix}_A = \begin{bmatrix} X\sigma + Y\rho \\ t^n \end{bmatrix}_A.$$

Hence

$$(\Phi(\omega))(f) = \Phi(f\omega)(1) = \Phi_{\sigma,\rho}(f\omega)(1) = (\Phi_{\sigma,\rho}(\omega))(f)$$

for any $\omega \in H_{MC_M}^1(C_M)$ and $f \in C_M$. That is, $\Phi = \Phi_{\sigma,\rho}$. $\square$

Now we fix a σ_0 and an invertible ρ_0 . All endomorphisms of the C_M -module $H_{MC_M}^1(C_M)$ are of the form $\Phi_{\sigma_0,\rho_0}^{-1} \circ \Phi_{\sigma,\rho}$. Since different pairs of σ and ρ determine different C_M -linear maps $\Phi_{\sigma,\rho}$, the completion $\widehat{C_M}$ of C_M can be described as the set $\{\Phi_{\sigma,\rho}\}_{\sigma,\rho \in \hat{A}}$ with addition

$$\Phi_{\sigma_1,\rho_1} + \Phi_{\sigma_2,\rho_2} = \Phi_{\sigma_1+\sigma_2,\rho_1+\rho_2},$$

unit Φ_{σ_0,ρ_0} , and multiplication

$$\Phi_{\sigma_1,\rho_1} * \Phi_{\sigma_2,\rho_2} = \Phi_{\sigma_1,\rho_1} \circ \Phi_{\sigma_0,\rho_0}^{-1} \circ \Phi_{\sigma_2,\rho_2}$$

given by composition of endomorphisms. If $\sigma_0 = 0$ and $\rho_0 = 1$, then

$$\Phi_{\sigma_1,\rho_1} * \Phi_{\sigma_2,\rho_2} = \Phi_{\sigma_1\rho_2+\sigma_2\rho_1-2\sigma_1\sigma_2t(z_0-a_0),\rho_1\rho_2-\sigma_1\sigma_2t^2(z_0-a_0)^2}.$$

Identify $\Phi_{\sigma,\rho}$ with $\rho + \sigma X$ in $\hat{A}[X]/(X + t(z - a_0))^2$, compare their additions and multiplications, we get the following description of $\widehat{C_M}$.

Corollary 7. $\widehat{C_M} \simeq \hat{A}[X]/(X + t(z - a_0))^2$

For $\rho \in C_M$, the endomorphism $\Phi_{0,1}^{-1} \circ \Phi_{0,\rho}$ of $H_{MC_M}^1(C_M)$ is multiplication by ρ . Therefore, with respect to the isomorphism in Corollary 7, the embedding $C_M \rightarrow \hat{A}[X]/(X + t(z - a_0))^2$ of completion is the composition

$$C_M \xrightarrow{\text{inclusion}} \hat{A} \xrightarrow{\text{canonical}} \hat{A}[X]/(X + t(z - a_0))^2.$$

REFERENCES

- [1] Y. Akizuki. Einige Bemerkungen über primäre Integritätsbereiche mit Teilerkettensatz. *Proc. Phys.-Math. Soc. Japan*, 17:327–336, 1935.
- [2] I-C. Huang. Pseudofunctors on modules with zero dimensional support. *Mem. Amer. Math. Soc.*, 114(548):xii+53, 1995.
- [3] I-C. Huang. Applications of residues to combinatorial identities. *Proc. Amer. Math. Soc.*, 125(4):1011–1017, 1997.
- [4] I-C. Huang. An explicit construction of residual complexes. *J. Algebra*, 225(2):698–739, 2000.
- [5] I-C. Huang. The residue theorem via an explicit construction of traces. *J. Algebra*, 245(1):310–354, 2001.
- [6] I-C. Huang. Residue methods in combinatorial analysis. In *Local Cohomology and its Applications*, volume 226 of *Lecture Notes in Pure and Appl. Math.*, pages 255–342. Marcel Dekker, 2001.
- [7] I. Kaplansky. *Commutative rings*. The University of Chicago Press, Chicago, Ill.-London, revised edition, 1974.
- [8] M. Reid. *Undergraduate commutative algebra*. Cambridge University Press, Cambridge, 1995.
- [9] R. Y. Sharp and H. Zakeri. Local cohomology and modules of generalized fractions. *Mathematika*, 29(2):296–306 (1983), 1982.

INSTITUTE OF MATHEMATICS, ACADEMIA SINICA, NANKANG, TAIPEI 11529, TAIWAN, R.O.C.
 AND INSTITUTE OF INFORMATION SCIENCE, ACADEMIA SINICA, NANKANG, TAIPEI 11529, TAIWAN,
 R.O.C.

E-mail address: ichuang@math.sinica.edu.tw and ljl@iis.sinica.edu.tw